

**AMENDMENTS TO THE CLAIMS**

**This listing of claims will replace all prior versions and listings of claims in the application:**

**LISTING OF CLAIMS:**

- 1-12. (canceled)
13. (withdrawn-previously presented): A method of preparing the *Mycobacterium* promoter of claim 23, said process comprising the steps of:
- (a) isolating said promoter from *Mycobacterium* DNA,
  - (b) ligating the isolated promoter sequence of step (a) into a plasmid vector.
14. (withdrawn-previously presented): The process of claim 13, wherein the *Mycobacterium* promoter is 2.5 fold more active in *M. Smegmatis* than the heat shock protein promoter (P<sub>hsp60</sub>).
15. (withdrawn-previously presented): A process of expressing a reporter gene in *M. smegmatis* under carbon starved conditions, the process comprising the step of growing *M. smegmatis* containing the promoter of claim 28, wherein the carbon source is about 2.5 to 0.001% glucose.
16. (withdrawn-previously presented): The process of claim 15, wherein the carbon source is about 2 to 0.02% glucose.

17. (withdrawn-previously presented): The process of claim 15, wherein the growth of the *M. smegmatis* is reduced by about 6 to 25% by the presence of ethambutol.

18. (withdrawn-previously presented): The process of claim 17, wherein the growth of the *M. smegmatis* is reduced by about 7 to 21% by the presence of ethambutol.

19. (withdrawn-previously presented): The process of claim 15, wherein the growth of the *M. smegmatis* is reduced by about 15 to 45% by the presence of isoniazid.

20. (withdrawn-previously presented): The process of claim 19, wherein the growth of the *M. smegmatis* is reduced by about 18 to 40 % in the presence of isoniazid.

21. (withdrawn-previously presented): The process of claim 15, wherein the growth of the *M. smegmatis* is reduced by about 20 to 45% by the presence of rifampicin.

22. (withdrawn-previously presented): The process of claim 21, wherein the growth of the *M. smegmatis* is reduced by about 21 to 41% by the presence of rifampicin.

23. (currently amended): A *Mycobacterium* promoter, wherein the promoter is stable in *M. smegmatis* and *E. coli*, and consists of SEQ ID NO:2, a 200 base pair fragment upstream and proximal to the start codon of the *Mycobacterium tuberculosis* *relA*/SpoT coding sequence.

24. (previously presented): The *Mycobacterium* promoter of claim 23, wherein the promoter is operatively linked to a reporter gene.

25. (previously presented): The *Mycobacterium* promoter of claim 24, wherein said reporter gene is LacZ.

26. (previously presented): The *Mycobacterium* promoter of claim 24, wherein said reporter gene is xyle.

27. (previously presented): The *Mycobacterium* promoter of claim 24, wherein the promoter is 2.5 fold more active in *M. smegmatis* than a heat shock protein 60 promoter, P<sub>hsp60</sub>.

28 (previously presented): The *Mycobacterium* promoter of claim 24, wherein the promoter is further contained in a plasmid with an Ampicillin or Kanamycin resistance marker.

29. (canceled).

30. (canceled).